

GROWERTALKS

Culture Notes

1/15/2009

Daniels Plant Food: Does it work?

Paul Nelson, Carl Niedziela, and Dharma Pitchay

CORRECTION: In the print edition, the last paragraph contained an incorrect reference to a 100:1 fertilizer proportioner. It should have read: "When a 500:1 fertilizer proportioner draws fertilizer concentrate straight from the Daniels container, the resulting final concentration is 250 ppm N." This online article has been updated to reflect the correction.

Daniels Plant Food, a complete soybean-based liquid fertilizer with a 10N:4P2O5:3K2O analysis manufactured and marketed by DP Foods, LLC of Sherman, Texas, was developed in 1982 by Ralph Daniels. Beginning in 1986, grower and university trials were conducted on cut flowers, perennials, woody ornamentals, bedding plants and potted flowering plants, as well as fruit and vegetable plants. Marketing of the product began in 1996. The American Association of Plant Food Control Officials (AAPFCO) recognized "oilseed extract," the base for Daniels Plant Food, as a source of plant nutrients in 1998.

Although not classified as an organic fertilizer, Daniels Plant Food does contain sufficient biodegradable carbon to give it organic properties. The base for this fertilizer is produced from soybean seed. After roasting and crushing, oils are extracted from the seeds to yield a crude oil. Additional oils are extracted using potassium hydroxide or sodium hydroxide to yield high-value cooking (edible) oil. The remaining material is divided into two fractions—solid material containing fiber and protein used for animal feed and oilseed extract, the water-soluble portion of the seed. It contains minerals and water-soluble compounds such as amino acids, organic acids and sugars. Unless converted to fertilizer, oilseed extract constitutes a disposal challenge requiring waste treatment to eliminate carbon and lower nitrogen (N) content and ultimate discharge of remaining minerals, particularly phosphorus, into the environment.

For fertilizer production purposes, potassium hydroxide, rather than sodium hydroxide, is used for oil extraction from the seeds and additional inorganic nutrients are added to the resulting oilseed extract to bring it to Daniels' guaranteed analysis. Its final N composition includes 3.70% ammoniacal-N (A-N), 1.90% nitrate-N, 3.65% urea-N and 0.75% organic-N for a total of 81% of the N in the reduced forms of ammonium, urea and organic N. This fertilizer could also be produced from other types of oilseeds such as sunflower, corn and canola. An estimated 325,000 metric tons of oilseed extract are available in the U.S. per year. In addition to relieving the environmental burden caused by discharge of nutrients from oilseed extract, less phosphate and potassium (K) minerals would be mined from world reserves and less natural gas burned to produce ammonia, the base N stock in fertilizers.

When testing this nutrient source, we posed three questions. Could its 10:4:3 analysis meet crops' nutritional requirements? Would it supply adequate K with its 10N:3K2O ratio? And, would this fertilizer, which contains 81% of its N in reduced forms, result in ammonium toxicity? Grower acceptance of Daniels suggested these issues were

not a problem, but scientific verification was lacking. We conducted three experiments using conventional production methods to examine our concerns.

Petunia

We grew petunia Dreams Midnight in bedding plant flats using three fertilizer formulations (Daniels, 40% A-N, and 70% A-N) at three constant fertilizer rates (50, 100 and 200 ppm). The 40% A-N fertilizer was the same as commercial 20-10-20, while the 70% A-N fertilizer differed from 20-10-20 only in its higher proportion of N (70%) in the ammoniacal form.

Although plant size increased with increasing Daniels fertilization rate, the visual quality rating was highest at 100 ppm N. At 200 ppm, plants became too large for the container space and etiolated stems did not adequately support them. Days to flower was lowest at 100 and 200 ppm N. Considering all of the variables, the best rate of application for Daniels fertilizer was 100 ppm N.

Plants fertilized at 50 ppm N were lighter green and smaller than desired regardless of fertilizer type. Within this 50 ppm N group, plants fertilized with Daniels were deeper green, taller, rated higher and flowered earlier than those fertilized with the other two sources. Within the 100 ppm N group, the Daniels fertilized plants flowered in 4.5% less time than plants fertilized with the 40% A-N formulation and rated higher than the other two treatments. Plants in the 200 ppm N group grew too large for the container, resulting in thin stems and a problem with plant toppling. No differences occurred in plant size or days to flower across fertilizer types. Ratings were best for Daniels and 70% A-N fertilized plants. Across all fertilizer types and rates, the most commercially desirable plants were those fertilized with 100 ppm N from Daniels fertilizer.

Cyclamen

We selected cyclamen as the test crop because it's reported to have one of the highest K requirements of greenhouse crops with a recommended 1N:2K2O ratio. We grew cyclamen Laser Rose in 5.5-in. pots, one plant per pot. We applied Daniels and the 40% A-N fertilizers with each irrigation. During the first six weeks, all fertilizers were applied at an N rate of 126 ppm. After six weeks, we increased the two fertilizers to an N rate of 175 ppm, as is customary in commercial practice. Growth in the forms of canopy height, flower height above the canopy, plant diameter, plant fresh weight, corm fresh weight, and numbers of shoots, leaves and buds did not differ between Daniels and 40% A-N fertilizer. For these parameters, Daniels fertilizer performance was equal to the commercial fertilizer (40% A-N).

Twelve weeks after potting, Daniels-fertilized plants had an average of 9.7 open flowers compared to 6.6 flowers for the 40% A-N plants. This was an increase of 47% attributed to Daniels. This increase would allow a grower to market a showier plant or market the plants about one week earlier than normal with the standard number of flowers.

Bedding plants

In the third experiment, we planted plugs of dianthus Floral Lace, pansy Crystal Bowl Orange, petunia Dreams Red, salvia Victoria Blue, snapdragon Bell Mix, verbena Quartz Mix, and vinca Grape Cooler in bedding plant flats at two pH levels (standard and low) established using the standard and one-third standard lime rates. We applied three fertilizers (Daniels, 25% A-N, and 75% A-N) during each irrigation at rates of 100 or 200 ppm N. Flowering was similar in the seven bedding plant species when fertilized with Daniels compared to the conventional fertilizers. Daniels fertilizer resulted in a deeper green color in all species except dianthus, where color was similar. Generally speaking, high ammonium 75% A-N-treated plants were more compact (shorter and less leaf area) in all species

except verbenas, where there was no difference from the low ammonium 25% A-N plants. Plants treated with Daniels were equivalent to 75% A-N-treated plants in all comparisons except height in petunia, where they were similar to the taller 25% A-N plants. The compact Daniels and 75% A-N plants were more desirable for the tight space constraints of bedding plant flats.

Potassium

The K content of Daniels was adequate to meet the requirements of the eight species tested across the three experiments, including cyclamen, which has one of the highest K requirements for greenhouse crops. No symptoms of K deficiency or growth suppression occurred in plants treated with Daniels. Although leaf tissue K concentrations were lower than in plants treated with the conventional formulations, K levels were still above the minimum concentrations reported by Gibson et al. 2008 (Nutrient Deficiencies in Bedding Plants, Ball Publishing).

Ammonium toxicity

There was no ammonia toxicity in the first two experiments. In the third experiment, Daniels did not cause ammonium toxicity in any of the bedding plant species when applied at normal fertilizer rates (100 ppm) using the standard limestone rate (10 lbs./cubic yd.). When the Daniels rate applied to the seven bedding plant species was doubled to 200 ppm, ammonium toxicity still did not develop. However, when the lime rate was decreased to one-third (3.3 lbs./cubic yd.), resulting in excessively low substrate pH levels, a slight ammonium toxicity occurred in only one species (pansy) at both Daniels rates (100 and 200 ppm). The low ammonium 25% A-N fertilizer at the low lime rate caused slight symptoms in pansy and petunia while the high ammonium 75% A-N fertilizer caused moderate symptoms in dianthus, petunia, salvia and snapdragon and heavy symptoms in pansy, verbenas and vinca. Although we anticipated applying Daniels under extreme conditions of high concentration and low substrate pH would lead to ammonium toxicity, for the most part this did not occur. This indicates that, unlike conventional fertilizers, Daniels is very resistant to ammonium toxicity.

pH buffering

The lack of ammonium toxicity from Daniels may be due to pH buffering capacity since high pH protects against ammonium toxicity. Daniels has 81% of the N in reduced form (ammonium, organic-N and urea), resulting in a potential acidity of 327. Consequently, we expected the pH in the Daniels treatments to be low as expected from the high ammoniacal fertilizers (70% and 75% A-N). Surprisingly, substrate pH in the Daniels treatments was buffered at a higher pH similar to or higher than that in the low ammoniacal fertilizers (25% and 40% A-N) in all but two species (verbenas and vinca).

Electrical conductivity (EC)

Substrate salt levels were 50% to 75% lower in plants treated with Daniels. Only when Daniels was applied at a N level double that of the conventional formulations, were the substrate salt levels equivalent. This is due to the high percentage of N in urea (36.5%) and organic N (7.5%). Growers who normally use substrate EC as a guide for fertilization will need to factor in the lower EC of the Daniels.

Nutrient uptake

When all experiments are taken into consideration, Daniels, compared to the 25% A-N and 40% A-N fertilizers, generally resulted in lower tissue concentrations of K, Ca, Mg, and Zn and a higher Fe concentration. Less frequently, P and Cu concentrations were found to be higher.

Summary

Daniels Plant Food proved to be an effective fertilizer for seven species of bedding plants and cyclamen. Plants

tended to be desirably more compact for most species while for the others, growth was similar to the conventional 20-10-20-fertilized plants. Plant color with Daniels was deeper green in all but two species. Earlier flowers formed in petunia and cyclamen plants. Tissue K levels were lower but in all cases adequate. Ammonium toxicity was not a problem with Daniels. It was more resistant to ammonium toxicity than conventional fertilizers. Substrate EC levels were lower with Daniels. Daniels also provided substrate pH buffering against decline.

Daniels Plant Food is a 10-4-3 liquid formulation. When a **500:1 fertilizer proportioner** draws fertilizer concentrate straight from the Daniels container, the resulting final concentration is 250 ppm N. Since this avoids any pre-dilution of the concentrate, the manufacturer recommends twice weekly applications of Daniels at 250 ppm N for bedding plants. Clear water should be used for irrigations between fertilizer applications.